

Second class particles as microscopic characteristics in totally asymmetric nearest-neighbor K -exclusion processes

2000

Timo Seppäläinen

Department of Mathematics
Iowa State University
Ames, Iowa 50011, USA
seppalai@iastate.edu

Summary. We study aspects of the hydrodynamics of one-dimensional totally asymmetric K -exclusion, building on the hydrodynamic limit of Seppäläinen (1999). We prove that the weak solution chosen by the particle system is the unique one with maximal current past any fixed location. A uniqueness result is needed because we can prove neither differentiability nor strict concavity of the flux function, so we cannot use the Lax-Oleinik formula or jump conditions to define entropy solutions. Next we prove laws of large numbers for a second class particle in K -exclusion. The macroscopic trajectories of second class particles are characteristics and shocks of the conservation law for the particle density. In particular, we extend to K -exclusion Ferrari's result that the second class particle follows a macroscopic shock in the Riemann solution. The technical novelty of the proofs is a variational representation for the position of a second class particle, in the context of the variational coupling method.

Mathematics Subject Classification: Primary 60K35, Secondary 82C22

Key words: Exclusion process, second class particle, characteristic, hydrodynamic limit, variational coupling method

Short Title: Second class particles in K -exclusion

1. INTRODUCTION

This paper continues the study of the hydrodynamics of totally asymmetric nearest-neighbor K -exclusion processes. These are interacting systems of indistinguishable particles on the one-dimensional integer lattice \mathbf{Z} , subject to the restriction that each site contain at most K particles. Independently at each site and at constant exponential rate one, one particle jumps to the next site on the right, provided the K -exclusion rule is not violated.

A hydrodynamic limit for this process was proved in Seppäläinen (1999). This is a law of large numbers according to which the empirical particle density converges to a weak solution of a nonlinear conservation law, under the appropriate scaling of space and time.

The present paper begins with the question of uniqueness of the weak solution chosen by the particle system. The criterion we give is maximization of current across any fixed position. This issue arises because we do not know whether the flux function of the conservation law is strictly concave or differentiable. Consequently we cannot use the Lax-Oleinik formula to define an entropy solution, or one-sided bounds on jumps as an entropy criterion.

We address this difficulty because it is closely tied to the main topic of the paper, the dynamics of a second class particle in the hydrodynamic scaling. As we shall see, our incomplete understanding of the macroscopic equation constrains the macroscopic description of the second class particle.

A *second class particle* is an extra particle in the system that yields right of way to the regular, first class particles. The second class particle can jump only if there is no first class particle at its site who can jump, and a second class particle has to jump backward if first class particles fill its site.

Second class particles in asymmetric exclusion processes have been studied for several reasons. In a certain technically precise sense, the second class particle marks the location of a shock at the microscopic particle level. These ideas were developed by Ferrari, Kipnis and Saada (1991), Ferrari (1992), and Ferrari and Fontes (1994a). At this same time, Ferrari (1992) discovered that in the hydrodynamic limit a second class particle follows the characteristics of the macroscopic equation. The connection between second class particles and characteristics was further elucidated by Rezakhanlou (1995) and Ferrari and Kipnis (1995). The second class particle also enters naturally in the study of the fluctuations of the current, as in Ferrari and Fontes (1994b).

Another strain of work, technically related to the investigation of microscopic shocks, concerns the invariant distributions of the exclusion process as seen from a second class particle, and the invariant distributions of the two-species process that has positive densities of both first and second class particles. This program culminated in the complete description of these measures by Ferrari, Fontes and Kohayakawa (1994). These and other results, such as the central limit behavior of the second class particle, are discussed in Part III of Liggett (1999) and in the notes of Chapter 9 in Kipnis and Landim (1999).

The above results are for systems whose invariant distributions are explicitly known, namely the exclusion and zero-range processes. And the proofs typically use precise and deep knowledge about those invariant measures.

In the present paper we prove laws of large numbers for a single second class particle in the hydrodynamic scaling of totally asymmetric K -exclusion. Essentially no

information is currently available about the invariant distributions of these asymmetric K -exclusion processes.

Of the papers mentioned above, Ferrari (1992) and Rezakhanlou (1995) are the closest predecessors of our paper. Both derived laws of large numbers for the second class particle. The limiting trajectory is a characteristic or a shock of the conservation law of the particle density. Ferrari proved strong laws in an asymmetric simple exclusion process, mainly in these settings: the initial particle distribution is either an i.i.d. product measure, or a product measure that represents a shock profile that is constant on either side of the origin (this is known as the Riemann problem). Ferrari's proofs rely on couplings of processes of several species of particles, and knowledge of invariant distributions.

Rezakhanlou (1995) proved L^1 -laws of large numbers for the second class particle in asymmetric simple exclusion and in a totally asymmetric zero-range process. He assumed that initial distributions are product measures, and the initial macroscopic profiles are bounded integrable functions on \mathbf{R} , with some additional technical assumptions. His proof derives the limit of a single second class particle from the limit of a macroscopically visible interval of second class particles, and the latter he gets from the basic hydrodynamic limits for exclusion and zero-range processes, from Rezakhanlou (1991). The invariant distributions of the processes enter his proofs through the macroscopic equations. These are known explicitly precisely because the invariant distributions are known. Rezakhanlou's approach would prove some result for K -exclusion, since one can use the hydrodynamic limit of Seppäläinen (1999) instead of Rezakhanlou (1991).

Our approach is different from the earlier ones. We use a representation of the second class particle in the variational coupling method. To preview this, let us switch from the exclusion process to the associated marching soldiers process (in other words, consider the interface process instead of the increments process). In this approach a process $z(i, t)$ from a general initial interface is coupled with auxiliary processes $w^k(i, t)$ that evolve from translated wedge-shaped initial profiles. The coupling is arranged so that $z(i, t)$ is the envelope of the auxiliary processes:

$$(1.1) \quad z(i, t) = \sup_{k \in \mathbf{Z}} w^k(i - k, t).$$

Let $X(t)$ denote the position of a second class particle added to the exclusion process $\eta(i, t) = z(i, t) - z(i - 1, t)$. The new tool is this variational expression:

$$(1.2) \quad X(t) = \inf\{i : z(i, t) = w^k(i - k, t) \text{ for some } k \geq X(0)\}.$$

Variational formula (1.1) is a microscopic analogue of the Hopf-Lax formula of viscosity solutions of Hamilton-Jacobi equations. In this same vein, (1.2) corresponds to the definition of forward characteristics through the Hopf-Lax formula. This justifies regarding the second class particle as a microscopic analogue of the characteristic, as expressed in the title of the paper.

Variational formulas (1.1)–(1.2) permit us to work directly on the paths of the process, and we do not need a particular form for the initial particle distributions. We need to assume that the initial particle profile and the initial position of the second class particle both converge under the appropriate space scaling. The result is a law of large numbers for the second class particle. The theorems give either convergence in probability or almost surely, depending on which kind is assumed

initially. The limit is a characteristic or a shock of the macroscopic partial differential equation.

Here we run into the difficulty mentioned above. For $2 \leq K < \infty$ (cases other than exclusion or zero-range) we do not have sufficient control of the regularity of the flux f of the conservation law. For this reason we cannot assert that macroscopically the second class particle follows the solution of $\dot{x} = f'(\rho(x, t))$, where $\rho(x, t)$ is the macroscopic particle density. Instead, we use the Hopf-Lax formula to define the macroscopic trajectory of the second class particle.

If the macroscopic characteristics are unique (there are no rarefaction fans) we do not need the assumption of an initial law of large numbers for the second class particle. In this case the second class particle follows, in the hydrodynamic scale, the macroscopically defined characteristic that emanates from its random initial position.

We improve earlier results in two ways: by broadening the class of processes beyond exclusion and zero-range, and by permitting more general initial distributions. In one important sense the results of Ferrari and Rezakhanlou remain more general than ours: their results cover an asymmetric simple exclusion process where particles can jump both right and left, while our results are restricted to *totally* asymmetric processes where particles jump only to the right. This is because a way to apply the variational coupling approach without total asymmetry has yet to be discovered.

Overview of the paper. In Section 2 we discuss the results. Section 2.1 reviews the hydrodynamics of totally asymmetric K -exclusion and zero-range processes (Theorem 1). We state a uniqueness criterion for the weak solution chosen by the particle system, valid for a concave flux that is not necessarily differentiable or strictly concave (Theorem 2). Section 2.2 states the laws of large numbers for the second class particle (Theorems 3 and 4). Section 2.3 comments on the various assumptions and conclusions.

In Section 2.4 we discuss a generalization of the K -exclusion and zero-range processes to which our results apply. In this model particles may jump in batches of varying sizes from one site to the next. Such a generalization is natural from a queueing perspective, where the particles are customers moving through a sequence of service stations.

Some standard tools of conservation laws are not at our disposal for lack of information about the regularity of the flux f of the conservation law. This problem is clarified in Section 3. Without strict concavity of f the Lax-Oleinik formula cannot be used, and the correct weak solution can have shocks prohibited by entropy criteria (Section 3.1). Without differentiability of f we lose some good properties of characteristics (Section 3.2). In section 3.3 we prove Theorem 2, the uniqueness result.

Section 4 reviews the essentials of the variational coupling approach to exclusion and zero-range processes from Seppäläinen (1998a,b,1999), and develops the formula for the second class particle. Section 5 proves the laws of large numbers for the second class particle. The proofs are an interplay of the particle level properties from Section 4 and the properties of the macroscopic characteristics.

Notation. The set of natural numbers is $\mathbf{N} = \{1, 2, 3, \dots\}$, and the set of non-negative integers $\mathbf{Z}_+ = \{0, 1, 2, 3, \dots\}$.

2. RESULTS

2.1. Hydrodynamic limits and entropy solutions. We start by reviewing the hydrodynamic limit of totally asymmetric K -exclusion. The parameter K is a fixed positive finite or infinite integer $K \in \mathbf{N} \cup \{\infty\}$. Each process consists of indistinguishable particles that move on the one-dimensional integer lattice \mathbf{Z} , subject to the restriction that a site $i \in \mathbf{Z}$ can be occupied by at most K particles. If $K = \infty$ this constraint is absent. The state space of the process is $\mathcal{S} = \{0, 1, 2, \dots, K\}^{\mathbf{Z}}$ for finite K , and $\mathcal{S} = \mathbf{Z}_+^{\mathbf{Z}}$ if $K = \infty$. A generic element of \mathcal{S} is $\eta = (\eta(i) : i \in \mathbf{Z})$, where $\eta(i)$ denotes the number of particles present at site i .

The dynamics is such that at exponential rate 1, one particle moves from site i to $i + 1$, provided there is at least one particle at i and at most $K - 1$ particles at $i + 1$. The jump attempts happen independently at all sites i .

To realize this dynamics, let $\{D_i : i \in \mathbf{Z}\}$ denote a collection of mutually independent rate 1 Poisson point processes on the time axis $(0, \infty)$. We usually think of D_i as the random set of jump times (or *epochs*, to borrow Feller's term) rather than as the corresponding random step function. At each epoch of D_i a jump from site i to site $i + 1$ is attempted, and successfully executed if the rules permit. The state of the process at time t is the configuration $\eta(t) = (\eta(i, t) : i \in \mathbf{Z})$ of occupation numbers.

The dynamics can be represented by the generator L that acts on bounded cylinder functions f on \mathcal{S} :

$$(2.1) \quad Lf(\eta) = \sum_{i \in \mathbf{Z}} \mathbf{1}\{\eta(i) \geq 1, \eta(i+1) \leq K-1\} [f(\eta^{i,i+1}) - f(\eta)].$$

Here $\eta^{i,i+1}$ is the configuration that results from the jump of a single particle from site i to site $i + 1$:

$$\eta^{i,i+1}(j) = \begin{cases} \eta(i) - 1, & j = i \\ \eta(i+1) + 1, & j = i+1 \\ \eta(j), & j \neq i, i+1. \end{cases}$$

Special cases. (a) When $K = 1$ this process is known as the totally asymmetric simple exclusion process (TASEP). The Bernoulli probability measures $\{\nu^\rho\}$ on \mathcal{S} , indexed by density $\rho \in [0, 1]$, are invariant for the process. These are defined by the condition

$$\nu^\rho \{\eta(i_1) = \eta(i_2) = \dots = \eta(i_m) = 1\} = \rho^m$$

for any finite set $\{i_1, \dots, i_m\}$ of sites.

(b) If $K = \infty$ this process is a member of the family of totally asymmetric zero-range processes (TAZRP). The i.i.d. geometric probability measures $\{\nu^\rho\}$, $\rho \in [0, \infty)$, are invariant. These are given by

$$(2.2) \quad \nu^\rho \{\eta(i_1) = k_1, \eta(i_2) = k_2, \dots, \eta(i_m) = k_m\} = \prod_{s=1}^m \frac{1}{1+\rho} \left(\frac{\rho}{1+\rho} \right)^{k_s}$$

for distinct sites i_1, \dots, i_m and arbitrary integers $k_1, \dots, k_m \in \mathbf{Z}_+$.

(c) For $2 \leq K < \infty$ we call this the totally asymmetric K -exclusion process. For this case the existence of spatially ergodic invariant measures for all densities has not been proved.

The basic result about the large-scale behavior of these processes is the following hydrodynamic scaling limit. Let ρ_0 be a bounded measurable function on \mathbf{R} such that $0 \leq \rho_0(x) \leq K$. Assume given a probability space on which are defined a sequence of initial configurations $(\eta_n(i, 0) : i \in \mathbf{Z})$, $n \in \mathbf{N}$, and the Poisson jump time processes $\{D_i\}$. The processes $\{\eta_n(\cdot)\}$ are constructed on this probability space by the familiar graphical method. Assume that at time zero the following strong law of large numbers is valid:

$$(2.3) \quad \text{For all } -\infty < a < b < \infty, \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=[na]+1}^{[nb]} \eta_n(i, 0) = \int_a^b \rho_0(x) dx \quad \text{a.s.}$$

Theorem 1. *Under assumption (2.3) a strong law of large numbers holds for all $t > 0$ and all $-\infty < a < b < \infty$:*

$$(2.4) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=[na]+1}^{[nb]} \eta_n(i, nt) = \int_a^b \rho(x, t) dx \quad \text{a.s.}$$

The deterministic limiting density $\rho(x, t)$ is a weak solution of the Cauchy problem

$$(2.5) \quad \rho_t + f(\rho)_x = 0, \quad \rho(x, 0) = \rho_0(x).$$

The flux function f of the conservation law (2.5) is a fixed, deterministic concave function on $[0, K]$ that depends on K .

The limit (2.3) is the only assumption needed in the following strong sense: The joint distribution of the initially defined random variables $\{(\eta_n(i, 0) : i \in \mathbf{Z}) : n \in \mathbf{N}\}$, and $\{D_i : i \in \mathbf{Z}\}$ may be arbitrary. As long as the marginal distribution of $\eta_n(i, 0)$ satisfies (2.3), and marginally $\{D_i\}$ are i.i.d. rate 1 Poisson point processes, Theorem 1 is valid.

A second point is that if (2.3) holds in probability, then the conclusion (2.5) also holds in probability.

For exclusion ($K = 1$) and zero-range ($K = \infty$) Theorem 1 has gone through many stages of generalization, beginning with the seminal work of Rost (1981). The entropy technique of Rezakhanlou (1991) proves it for multidimensional exclusion and zero-range processes, without requiring total asymmetry. The variational coupling method of Seppäläinen (1998a,b) permits the most general initial distributions for the one-dimensional, totally asymmetric case. For $2 \leq K < \infty$ Theorem 1 was first proved in Seppäläinen (1999).

Let us write f_K for the flux function f in (2.5) when we emphasize dependence on K . f_K is known in cases where invariant distributions are explicitly known: for exclusion $f_1(\rho) = \rho(1 - \rho)$ for $0 \leq \rho \leq 1$, and for zero-range $f_\infty(\rho) = \rho/(1 + \rho)$ for $0 \leq \rho < \infty$. For $1 < K < \infty$ we know the concavity and continuity of f_K , the symmetry $f_K(\rho) = f_K(K - \rho)$, the monotonicity $f_K(\rho) \leq f_{K+1}(\rho)$, and these bounds:

$$(2.6) \quad \min\{\rho(1 - \rho), 1/4\} \leq f_K(\rho) \leq \frac{\rho}{K} \quad \text{for } 0 \leq \rho \leq K/2.$$

Currently we have no information about the regularity of f for $1 < K < \infty$. As suggested by f_1 and f_∞ , we would expect f to be C^1 and strictly concave, in other words, for its graph to have no corners or line segments. This lack of information about f creates a problem because the common definitions of entropy solutions of (2.5) are valid only for strictly convex or concave flux functions. Thus we cannot apply standard entropy criteria such as one-sided bounds on jumps or the Lax-Oleinik formula. We show this in Section 3.

The next task is to complement Theorem 1 with a uniqueness criterion. First we explain how the solution $\rho(x, t)$ that the particle system chooses in Theorem 1 is defined. In the next discussion assume first $K < \infty$. Afterwards we comment on the $K = \infty$ case.

Define an antiderivative u_0 of ρ_0 by

$$(2.7) \quad u_0(0) = 0 \text{ and } u_0(b) - u_0(a) = \int_a^b \rho_0(x) dx \text{ for all } a < b.$$

Let g be the nonincreasing, nonnegative convex function on \mathbf{R} defined by

$$(2.8) \quad g(x) = \sup_{0 \leq \rho \leq K} \{f(\rho) - x\rho\}.$$

Symmetry of f and bounds (2.6) imply that $f'(0) = 1$ and $f'(K) = -1$, the existence of the derivatives being part of the conclusion. It follows that g satisfies

$$(2.9) \quad g(x) = 0 \text{ for } x \geq 1 \text{ and } g(x) = -Kx \text{ for } x \leq -1.$$

Then for $x \in \mathbf{R}$ set $u(x, 0) = u_0(x)$, and for $t > 0$

$$(2.10) \quad u(x, t) = \sup_{y \in \mathbf{R}} \left\{ u_0(y) - tg\left(\frac{x-y}{t}\right) \right\}.$$

The supremum is attained at some $y \in [x-t, x+t]$. The function u is uniformly Lipschitz on $\mathbf{R} \times [0, \infty)$, nonincreasing in t and nondecreasing in x . (2.10) is known as the *Hopf-Lax formula*. It defines $u(x, t)$ as a *viscosity solution* of the Hamilton-Jacobi equation

$$(2.11) \quad u_t + f(u_x) = 0, \quad u(x, 0) = u_0(x)$$

[Ch. 10 in Evans (1998)]. The viscosity solution is unique by Theorem 2.1 of Ishii (1984) that applies to (2.11) because f is continuous and u is uniformly continuous.

If $K = \infty$ we know that $g(x) = (1 - \sqrt{x})^2$ for $0 \leq x \leq 1$, $g \equiv 0$ on $(1, \infty)$ and we can take $g \equiv \infty$ on $(-\infty, 0)$. The supremum in (2.10) is attained at some $y \in [x-t, x]$. Since ρ_0 is assumed bounded, u is again Lipschitz on $\mathbf{R} \times [0, \infty)$.

Let

$$(2.12) \quad \rho(x, t) = \frac{\partial}{\partial x} u(x, t)$$

be the a.e. defined x -derivative. Recall this standard definition: a locally bounded measurable function $\lambda(x, t)$ is a *weak solution* of (2.5) if for all compactly supported and continuously differentiable test functions ϕ on $\mathbf{R} \times [0, \infty)$,

$$(2.13) \quad \int_0^\infty \int [\lambda(x, t)\phi_t(x, t) + f(\lambda(x, t))\phi_x(x, t)] dx dt + \int \rho_0(x)\phi(x, 0) dx = 0.$$

By the proof of Thm. 3.4.2 in Evans (1998), $\rho(x, t)$ is a weak solution of (2.5) in the integral sense (2.13). The solution $\rho(x, t)$ defined by (2.12) is the one chosen by the particle system in Theorem 1.

In cases $K = 1$ and $K = \infty$ we know that the flux f is differentiable and strictly concave. Then $\rho(x, t)$ can be equivalently defined by the Lax-Oleinik formula, and we can call $\rho(x, t)$ the “unique entropy solution.” For $2 \leq K < \infty$ we can only define $\rho(x, t)$ through (2.12). In Sections 3.1–3.2 we show that without further knowledge about f this solution cannot be defined by the Lax-Oleinik formula, and we cannot rule out discontinuities of the wrong kind that violate entropy conditions.

To complement the characterization of $\rho(x, t)$ through the viscosity solution of (2.11), we give a uniqueness criterion for which strict concavity or differentiability of f are not needed.

Theorem 2. *Among the weak solutions of (2.5), $\rho(x, t)$ of (2.12) is characterized by maximal current over time. More precisely, suppose $\lambda(x, t)$ is a nonnegative, locally bounded measurable function on $\mathbf{R} \times (0, \infty)$ that satisfies the integral criterion (2.13). Fix $t > 0$. Then for almost all $x \in \mathbf{R}$,*

$$(2.14) \quad \int_0^t f(\lambda(x, s)) ds \leq \int_0^t f(\rho(x, s)) ds.$$

Conversely, if equality holds in (2.14) for almost all (x, t) , then $\lambda(x, t) = \rho(x, t)$ for almost all (x, t) .

Thus the particle system chooses the weak solution that transports the most material. Theorem 2 is proved in Section 3.3. The system maximizes the current because f is concave. With a convex flux the current would be minimized, as is the case with the stick model version of Hammersley’s process, see Seppäläinen (1996).

2.2. Second class particles and characteristics. In this section we state the laws of large numbers for the second class particle. The representation of the second class particle in the variational coupling is in Section 4.2.

Let $X(t)$ denote the position of a single second class particle added to the process $\eta(t)$. $X(t)$ is rigorously defined as follows. Assume given a probability space on which are defined the following random variables: the Poisson jump time processes $\{D_i\}$, an initial particle configuration $(\eta(i, 0) : i \in \mathbf{Z})$, and an initial location $X(0)$. Define a second initial configuration $(\tilde{\eta}(i, 0) : i \in \mathbf{Z})$ that differs from η only at site $X(0)$: $\tilde{\eta}(X(0), 0) = \eta(X(0), 0) + 1$ and $\tilde{\eta}(i, 0) = \eta(i, 0)$ for $i \neq X(0)$. [Of course we must have $\eta(X(0), 0) \leq K - 1$ a.s. for this to be possible.]

Run the processes $\eta(t)$ and $\tilde{\eta}(t)$ so that they read the same Poisson jump time processes $\{D_i\}$. This is known as the basic coupling of $\eta(t)$ and $\tilde{\eta}(t)$. There is always a unique site $X(t)$ at which the two processes differ: $\tilde{\eta}(X(t), t) = \eta(X(t), t) + 1$ and $\tilde{\eta}(i, t) = \eta(i, t)$ for $i \neq X(t)$. This defines the position $X(t)$ of the second class particle.

The way in which the second class particle yields to first class particles becomes evident when we consider the two types of transitions that can happen to it. Let $X(\tau-) = i$.

(i) Suppose τ is a jump epoch for Poisson process D_i . The second class particle can jump from i to $i+1$ if there is space at site $i+1$, but only if there is no first class particle to jump. In other words, if $\eta(i, \tau-) = 0$ and $\eta(i+1, \tau-) < K$, then $X(\tau) = i+1$. Otherwise the second class particle does not move and $X(\tau) = X(\tau-) = i$.

(ii) Suppose τ is a jump epoch for D_{i-1} . The second class particle jumps from i to $i - 1$ if first class particles fill site i . In other words, if $\eta(i - 1, \tau-) \geq 1$ and $\eta(i, \tau-) = K - 1$, then $X(\tau) = i - 1$.

If $K = \infty$ the second class particle never jumps left and only the first kind of transition happens to it.

The definition in terms of η and $\tilde{\eta}$ generalizes naturally to a two-species process of first and second class particles. But in this paper we consider only a single second class particle added to the process.

Next we discuss the characteristics of the conservation law (2.5). The lack of information about the regularity of f becomes a problem again. The well-known description of characteristics as generalized solutions of the ordinary differential equation $dx/dt = f'(\rho(x, t))$ [see Dafermos (1977)] does not work now. As Reza-khanlou (1995), we use the Hopf-Lax formula to define characteristics.

Once $u(x, t)$ is defined by (2.10), a semigroup property is in force: for all $0 \leq s < t$ and $x \in \mathbf{R}$,

$$(2.15) \quad u(x, t) = \sup_{y \in \mathbf{R}} \left\{ u(y, s) - (t - s)g\left(\frac{x - y}{t - s}\right) \right\}.$$

The supremum is assumed at some point $y \in [x - (t - s), x + (t - s)]$. (If $K = \infty$ this range would actually be $[x - (t - s), x]$.) By continuity the supremum in (2.15) is attained at the least and largest maximizers, denoted by

$$(2.16) \quad y^-(x; s, t) = \inf \left\{ y \geq x - (t - s) : u(x, t) = u(y, s) - (t - s)g\left(\frac{x - y}{t - s}\right) \right\}$$

and

$$(2.17) \quad y^+(x; s, t) = \sup \left\{ y \leq x + (t - s) : u(x, t) = u(y, s) - (t - s)g\left(\frac{x - y}{t - s}\right) \right\}.$$

Abbreviate $y^\pm(x, t) = y^\pm(x; 0, t)$. As functions of x for fixed $0 \leq s < t$, $y^+(x; s, t)$ is right-continuous, $y^-(x; s, t)$ is left-continuous, both are nondecreasing, and trivially from the definition $\lim_{x \rightarrow -\infty} y^\pm(x; s, t) = -\infty$ and $\lim_{x \rightarrow \infty} y^\pm(x; s, t) = \infty$. See comment 2.3.1 below for justification of the restriction $x - (t - s) \leq y \leq x + (t - s)$ in the definitions of $y^\pm(x; s, t)$.

The minimal and maximal forward characteristics are defined for $b \in \mathbf{R}$, $0 \leq s < t$, as

$$(2.18) \quad x^-(b; s, t) = \inf \{x : y^+(x; s, t) \geq b\}$$

and

$$(2.19) \quad x^+(b; s, t) = \sup \{x : y^-(x; s, t) \leq b\}.$$

Immediate properties are these:

As functions of b , $x^+(b; s, t)$ is right-continuous, $x^-(b; s, t)$ is left-continuous, and both are nondecreasing. For the $s = 0$ case abbreviate $x^\pm(b, t) = x^\pm(b; 0, t)$. As functions of t , $x^\pm(b; s, t)$ are Lipschitz and satisfy the initial condition $x^\pm(b; s, s) = b$. When characteristics are unique we write $x(b; s, t) = x^\pm(b; s, t)$.

Now consider the setting of the hydrodynamic limit Theorem 1, with a sequence of processes η_n that satisfy the initial limit (2.3). Let $X_n(t)$ denote the position of a second class particle in the process η_n . The initial position $X_n(0)$ may be deterministic or random, and may depend in an arbitrary fashion on the initial particle configurations $(\eta_n(i, 0))$ and the Poisson processes $\{D_i\}$. For the first result assume a law of large numbers at time zero: for a deterministic point $b \in \mathbf{R}$

$$(2.20) \quad \lim_{n \rightarrow \infty} \frac{X_n(0)}{n} = b \quad \text{a.s.}$$

Theorem 3. *Under assumptions (2.3) and (2.20), for any fixed $t > 0$,*

$$(2.21) \quad x^-(b, t) \leq \liminf_{n \rightarrow \infty} \frac{X_n(nt)}{n} \leq \limsup_{n \rightarrow \infty} \frac{X_n(nt)}{n} \leq x^+(b, t)$$

almost surely. In particular, if $x(b, t) = x^\pm(b, t)$, then we have a strong law of large numbers:

$$(2.22) \quad \lim_{n \rightarrow \infty} \frac{X_n(nt)}{n} = x(b, t) \quad \text{a.s.}$$

In the special case where characteristics are unique we can dispense with assumption (2.20). The second class particle tracks the macroscopically defined characteristic that emanates from its random initial position, as stated in part (a) of the next theorem. We only need to assume that the fluctuations of the initial location are macroscopically bounded as n grows:

$$(2.23) \quad -\infty < \liminf_{n \rightarrow \infty} \frac{X_n(0)}{n} \leq \limsup_{n \rightarrow \infty} \frac{X_n(0)}{n} < \infty \quad \text{almost surely.}$$

The uniqueness of characteristics is true if we look at characteristics started at a time $s > 0$. But only if the flux function is differentiable, so this assumption needs to be made explicitly for $2 \leq K < \infty$ in part (b).

Theorem 4. *Assume (2.3) and (2.23).*

(a) *If $x(y, t) = x^\pm(y, t)$ for all $y \in \mathbf{R}$, then almost surely*

$$(2.24) \quad \lim_{n \rightarrow \infty} \left| \frac{X_n(nt)}{n} - x\left(\frac{X_n(0)}{n}, t\right) \right| = 0.$$

(b) *Let $0 < s < t$. In the case $2 \leq K < \infty$ suppose f is differentiable on $[0, K]$. For $K = 1$ and $K = \infty$ this is known and does not need to be assumed. Then $x(y; s, t) = x^\pm(y; s, t)$ for all $y \in \mathbf{R}$, and almost surely*

$$(2.25) \quad \lim_{n \rightarrow \infty} \left| \frac{X_n(nt)}{n} - x\left(\frac{X_n(ns)}{n}; s, t\right) \right| = 0.$$

The meaning of $x(n^{-1}X_n(0), t)$ is that the deterministic function $x(\cdot, t)$ defined by (2.18–19) is evaluated at the random point $n^{-1}X_n(0)$. If assumption (2.23) is strengthened to $na \leq X_n(0) \leq nb$ for all large enough n , then for Theorem 4(a) it is sufficient to assume $x(y, t) = x^\pm(y, t)$ for $a \leq y \leq b$. As for Theorem 1, if the assumptions are valid in probability, then so are the conclusions.

2.3. Comments on the assumptions and results.

2.3.1. Differentiability and strict convexity. Since g is convex, left and right derivatives

$$g'_-(x) = \lim_{\varepsilon \searrow 0} \frac{g(x - \varepsilon) - g(x)}{-\varepsilon} \quad \text{and} \quad g'_+(x) = \lim_{\varepsilon \searrow 0} \frac{g(x + \varepsilon) - g(x)}{\varepsilon}$$

exist at all points x . Same is true for f , and for both f and g differentiability on an open interval is equivalent to continuous differentiability. *Strict convexity* means that there are no $x_1 < x_2$ such that $g'_+(x_1) = g'_-(x_2)$, which is the same as not having a line segment in the graph of g . From the duality (2.8) it follows that f is differentiable on $[0, K]$ iff g is strictly convex on $[-1, 1]$, and f is strictly concave on $[0, K]$ iff g is differentiable on \mathbf{R} . Consult Rockafellar (1970) for a general treatment of convex analysis.

Theorem 4(b) is true because strict convexity of g guarantees that $x(y; s, t) = x^\pm(y; s, t)$ for all $y \in \mathbf{R}$, if $s > 0$. This is proved in Proposition 3.1 in Section 3.2.

The restriction $x - (t - s) \leq y \leq x + (t - s)$ in definition (2.16–17) of $y^\pm(x; s, t)$ is necessary because g is linear outside $[-1, 1]$. For example, take as initial profile $\rho_0(x) = K$ for $x \leq 0$ and $\rho_0(x) = 0$ for $x > 0$. Then for $x > t$, the entire interval $[0, x - t]$ consists of maximizers in the Hopf-Lax formula. Without the restriction in (2.16) we would have $y^-(x, t) = 0$ for all $x > t$, and then $x^+(0, t) = \infty$ which is not the appropriate characteristic.

2.3.2. Shocks. Consider the simplest shock case (Riemann problem) with initial profile $\rho_0(x) = \alpha \mathbf{1}\{x < 0\} + \beta \mathbf{1}\{x > 0\}$ where $\alpha < \beta$. The solution (2.12) is $\rho(x, t) = \alpha \mathbf{1}\{x < \xi t\} + \beta \mathbf{1}\{x > \xi t\}$ with shock speed $\xi = (f(\beta) - f(\alpha))/(\beta - \alpha)$. The unique characteristic from the origin is $x(0, t) = \xi t$. Theorem 3 says that the second class particle converges to the shock. This generalizes results of Ferrari (1992) in two ways: From exclusion to K -exclusion, and from initial product distributions to general initial distributions that have the shock profile.

2.3.3. Rarefaction fan. Ferrari and Kipnis (1995) proved convergence in probability in (2.25) for the following case of TASEP: initial distribution is product measure with density ρ to the left and density λ to the right of the origin, with $\rho > \lambda$, and the second class particle initially at the origin. This initial profile $\rho_0(x) = \rho \mathbf{1}\{x < 0\} + \lambda \mathbf{1}\{x > 0\}$ has an inadmissible shock at the origin, which produces a “rarefaction fan” of characteristics with $x^-(0, t) = (1 - 2\rho)t$ and $x^+(0, t) = (1 - 2\lambda)t$. Ferrari and Kipnis also proved that weakly $n^{-1}X_n(nt)$ converges to the uniform distribution on $[x^-(0, t), x^+(0, t)]$. Consequently one cannot hope for a law of large numbers when $x^-(0, t) < x^+(0, t)$.

If f is not differentiable at ρ , then the situation $x^-(b, t) < x^+(b, t)$ happens even for the constant profile $\rho(x, t) \equiv \rho$. See Example 3.1 in Section 3.2. In this case we cannot deduce convergence of $n^{-1}X_n(nt)$ from Theorem 3, only the bounds (2.24), unless we can prove the differentiability of f .

2.3.4. The characteristic ordinary differential equation. When f is strictly concave and differentiable, Theorem 3 also contains this statement: The characteristics

$x(t) = x^\pm(b, t)$ are *Filippov solutions* of the o.d.e.

$$(2.26) \quad \frac{dx}{dt} = f'(\rho(x, t)), \quad x(0) = b,$$

where $\rho(x, t)$ is the Lax-Oleinik solution of (2.5). Since $\rho(x, t)$ does not exist at all x , (2.26) is interpreted as

$$(2.27) \quad \operatorname{ess\,inf}_x f'(\rho(x, t)) \leq \frac{dx}{dt} \leq \operatorname{ess\,sup}_x f'(\rho(x, t)) \quad \text{for a.e. } t.$$

The essential infimum relative to x , at the point (x, t) , is defined by

$$\operatorname{ess\,inf}_x f'(\rho(x, t)) = \lim_{\delta \searrow 0} \operatorname{ess\,inf}_{x-\delta < y < x+\delta} f'(\rho(y, t)),$$

and similarly for the essential supremum. The Lax-Oleinik formula (we review it in Section 3.1 below) shows that (2.27) is equivalent to

$$(2.28) \quad f' \left(-g' \left(\frac{x - y^+(x, t)}{t} \right) \right) \leq \frac{dx}{dt} \leq f' \left(-g' \left(\frac{x - y^-(x, t)}{t} \right) \right)$$

for almost every t . These results are developed in Rezakhanlou (1995) for the exclusion and zero-range processes.

At present we cannot ascertain strict concavity or differentiability of f for $2 \leq K < \infty$. This is a problem, because to interpret (2.26) in the Filippov sense, $f'(\rho(x, t))$ must exist at least a.e. But if f' does not exist at ρ and we consider the constant solution $\rho(x, t) \equiv \rho$, then $f'(\rho(x, t))$ does not exist at any (x, t) .

Without any assumptions on f beyond concavity, bounds (2.28) are valid for $x(t) = x^\pm(b, t)$ in the form

$$(2.29) \quad f'_+ \left(-g'_+ \left(\frac{x - y^+(x, t)}{t} \right) \right) \leq \frac{dx}{dt} \leq f'_- \left(-g'_- \left(\frac{x - y^-(x, t)}{t} \right) \right)$$

for almost every t . If we assume f differentiable, then in fact (2.26) is valid again in the sense (2.27), with $\rho(x, t)$ given by (2.12). In other words, the o.d.e. works even though the Lax-Oleinik formula fails without strict concavity of f . We omit the proof of (2.29) and the last statement above.

2.4. Generalization: particles jump in batches. We discuss here a generalization of the K -exclusion and zero-range processes for which the variational coupling technique works and Theorems 1–4 are true.

From a queueing perspective, the particles are customers moving through an infinite sequence of servers. Each server has space for K customers in its queue, and $\eta(i, t)$ is the number of customers waiting at server i at time t . In the formulation (2.1), each server serves at exponential rate 1, and only one customer is served at a time and moves on to the next server. We generalize this by permitting customers to move in batches of varying sizes, with exponential rates and independently for each batch size. Let β_h be the rate at which a batch of h customers can jump from i to $i+1$, $h = 1, 2, 3, \dots$. If the state just before the jump does not permit h

customers to jump [so either $\eta(i) < h$ or $\eta(i+1) > K - h$], we do not suppress the entire jump but instead move as many customers as possible without violating the constraints.

The family of processes is parametrized by K and a sequence of nonnegative numbers $(\beta_h : h \in \mathbf{N})$. If $K < \infty$ assume $\beta_h = 0$ for $h > K$. To realize this dynamics take a collection of mutually independent Poisson point processes $\{D_i^h : i \in \mathbf{Z}, h \in \mathbf{N}\}$ on the time axis $(0, \infty)$, so that the rate of D_i^h is β_h . At each epoch of D_i^h , a batch of h particles (customers) attempt to jump from site i to site $i+1$. The generator becomes

$$Lf(\eta) = \sum_{i \in \mathbf{Z}} \sum_{h \in \mathbf{N}} \beta_h [f(\eta^{h,i,i+1}) - f(\eta)],$$

where $\eta^{h,i,i+1}$ is the configuration that results from an attempt to move a batch of h particles from site i to $i+1$. Denote the actual number of particles that move by

$$b = b(h, \eta(i), \eta(i+1)) = \min\{h, \eta(i), K - \eta(i+1)\}.$$

Then the new configuration is

$$\eta^{h,i,i+1}(j) = \begin{cases} \eta(i) - b, & j = i \\ \eta(i+1) + b, & j = i+1 \\ \eta(j), & j \neq i, i+1. \end{cases}$$

The definition of a second class particle is the same as before. From the queueing perspective it is a second class customer. He can move forward only if first class customers do not fill the batch that is served, and he has to move back to the previous server if arriving first class customers fill the capacity of the queue.

The hydrodynamic limit Theorem 1 is valid as stated for the batch process too, provided the jump rates are restricted so that we can expect finite limits. It is enough to require $\sum_{h \in \mathbf{N}} \beta_h h^2 < \infty$. Under this assumption a weaker version of Theorem 1 with convergence in probability was proved in Seppäläinen (2000). That paper also shows how to adapt the variational coupling approach to batch jumps.

If $K = \infty$, the i.i.d. geometric probability measures (2.2) are again invariant, for any choice of the rates (β_h) . In this case the flux function is

$$f(\rho) = \sum_{h \in \mathbf{N}} \beta_h \left[\frac{\rho}{1+\rho} + \left(\frac{\rho}{1+\rho} \right) + \cdots + \left(\frac{\rho}{1+\rho} \right)^h \right].$$

3. SCALAR CONSERVATION LAWS WITH CONCAVE FLUX FUNCTIONS

In this section we investigate the consequences of the lack of strict concavity and differentiability of the flux function. It is important to clarify these facts because they directly influence the results we obtain for the particle models. Section 3.1 looks at strict concavity and section 3.2 at differentiability. In section 3.3 we prove the uniqueness criterion of Theorem 2 which assumes neither strict concavity nor differentiability of the flux.

3.1. Strict concavity of the flux and entropy shocks. We begin by recalling the Lax-Oleinik formula from Section 3.4 in Evans (1998), and then show how it depends on the strict concavity of the flux function.

Lax-Oleinik formula. Suppose f is strictly concave and differentiable on its domain $[0, K]$. Define two functions

$$(3.1) \quad \rho^\pm(x, t) = -g' \left(\frac{x - y^\pm(x, t)}{t} \right).$$

For fixed $t > 0$, under these assumptions $y^+(x, t) = y^-(x, t)$ for all but countably many x , and consequently the Lax-Oleinik solution

$$(3.2) \quad \rho_{\text{LO}}(x, t) = \rho^\pm(x, t)$$

is well-defined for all but countably many x . It satisfies $\rho_{\text{LO}}(x, t) = u_x(x, t)$ a.e., and is a weak solution of (2.5) in the integral sense (2.13).

In particular, when it exists ρ_{LO} is the solution defined in (2.12) for the hydrodynamic limit. One can check that $\rho_{\text{LO}}(\cdot, t)$ is continuous at all but the countably many x at which $\rho^-(x, t) \neq \rho^+(x, t)$. If $\rho_{\text{LO}}(\cdot, t)$ is discontinuous at x , there must be a jump upward: $\rho_{\text{LO}}(x-, t) < \rho_{\text{LO}}(x+, t)$. This is because g' is nondecreasing and $y^-(x, t) \leq y^+(x, t)$. Let us call such a jump an *entropy shock* for a concave flux function. [For a convex flux entropy shocks are downward jumps.]

There are two ways definition (3.1)–(3.2) could conceivably run into a problem:

(i) If f is not strictly concave then g' fails to exist at some points and the right-hand side of (3.1) may not be defined.

(ii) If f' does not exist everywhere then $y^-(x, t) < y^+(x, t)$ may happen for a nontrivial interval of points x , and formula (3.2) is not true a.e. See Example 3.1 in Section 3.2.

It turns out that if f is strictly concave, definition (3.1–2) works even if $y^-(x, t) = y^+(x, t)$ fails. And all discontinuities are entropy shocks.

Theorem 5. *Suppose the flux f is strictly concave. Then for a fixed $t > 0$ the solution $\rho(x, t) = u_x(x, t)$ defined as in (2.12) exists and is continuous on a set $H \subseteq \mathbf{R}$ that contains all but countably many x . On H $\rho(x, t) = \rho^\pm(x, t)$ so the Lax-Oleinik formula is valid. For all x $\rho(x-, t) \leq \rho(x+, t)$, so the discontinuities of $\rho(\cdot, t)$ are entropy shocks.*

Conversely, suppose f has a linear segment. Then there is a range of densities ρ such that the constant solution $\rho(x, t) \equiv \rho$ cannot be represented by any left or right derivative of g , so the Lax-Oleinik formula (3.1–2) fails. Furthermore, the relevant solution defined by (2.12) can have nonentropy shocks $\rho(x-, t) > \rho(x+, t)$.

Proof. Strict concavity of f implies that g is differentiable. Inequalities (3.3)–(3.4) below then imply that the right and left x -derivatives of $u(x, t)$ exist and are given by

$$u_{x\pm}(x, t) = -g' \left(\frac{x - y^\pm(x, t)}{t} \right) = \rho^\pm(x, t).$$

Since g' is continuous and $y^+(\cdot, t)$ nondecreasing and right-continuous, we see that $\rho^+(\cdot, t)$ is right-continuous, has at most countably many discontinuities, and has left limits at all points. And correspondingly for $\rho^-(x, t)$.

By the Lipschitz property $u(x, t)$ is differentiable at a.e. x and is the integral of its derivative. Thus

$$u(x, t) = \int_0^x u_x(\xi, t) d\xi = \int_0^x \rho^\pm(\xi, t) d\xi.$$

Consequently $\rho(x, t) \equiv u_x(x, t)$ exists and equals $\rho^\pm(x, t)$ at any x where one of $\rho^\pm(x, t)$ is continuous. This defines the set H . From this follows $\rho(x-, t) = \rho^-(x, t) \leq \rho^+(x, t) = \rho(x+, t)$.

For the converse, suppose $f'(\rho) = \xi$ for $\rho_0 < \rho < \rho_1$. Then g has a corner at ξ , with $g'_-(\xi) = -\rho_1$ and $g'_+(\xi) = -\rho_0$. The values $\rho \in (\rho_0, \rho_1)$ are not taken by g'_\pm at any point, so a constant solution $\rho(x, t) \equiv \rho \in (\rho_0, \rho_1)$ cannot be represented by a derivative of g .

The initial profile $\rho_0(x) = K$ for $x \leq 0$ and $\rho_0(x) = 0$ for $x > 0$ gives the solution $u(x, t) = -tg(x/t)$. Then $\rho((t\xi)-, t) = \rho_1 > \rho_0 = \rho((t\xi)+, t)$ is a nonentropy shock. \square

To finish this section we prove the inequalities used in the proof above. Let us write $\overline{\lim}_{\varepsilon \searrow 0}$ to simultaneously include both the lim sup and the lim inf as $\varepsilon \searrow 0$.

Lemma 3.1. *The following inequalities hold for right and left x -derivatives of $u(x, t)$ at fixed $t > 0$.*

$$(3.3) \quad -g'_+ \left(\frac{x - y^-(x, t)}{t} \right) \leq \overline{\lim}_{\varepsilon \searrow 0} \frac{u(x - \varepsilon, t) - u(x, t)}{-\varepsilon} \leq -g'_- \left(\frac{x - y^-(x, t)}{t} \right)$$

and

$$(3.4) \quad -g'_+ \left(\frac{x - y^+(x, t)}{t} \right) \leq \overline{\lim}_{\varepsilon \searrow 0} \frac{u(x + \varepsilon, t) - u(x, t)}{\varepsilon} \leq -g'_- \left(\frac{x - y^+(x, t)}{t} \right).$$

Proof. The proofs use convexity, the Hopf-Lax formula, and the right [left] continuity of $y^+(\cdot, t)$ and g'_+ [$y^-(\cdot, t)$ and g'_-]. As an illustration, here is the argument for the second inequality of (3.4). We leave the rest to the reader.

$$\begin{aligned} & u(x + \varepsilon, t) - u(x, t) \\ & \leq u_0(y^+(x + \varepsilon, t)) - tg \left(\frac{x + \varepsilon - y^+(x + \varepsilon, t)}{t} \right) \\ & \quad - u_0(y^+(x, t)) + tg \left(\frac{x - y^+(x, t)}{t} \right) \\ & = -\varepsilon \cdot \left(\frac{t}{\varepsilon} \right) \cdot \left\{ g \left(\frac{x - y^+(x + \varepsilon, t)}{t} + \frac{\varepsilon}{t} \right) - g \left(\frac{x - y^+(x, t)}{t} \right) \right\} \\ & \leq -\varepsilon \cdot g'_+ \left(\frac{x - y^+(x + \varepsilon, t)}{t} \right). \end{aligned}$$

Consider two possibilities: If $y^+(x + \varepsilon, t) = y^+(x, t)$ for small enough $\varepsilon > 0$, then

$$g'_+ \left(\frac{x - y^+(x + \varepsilon, t)}{t} \right) = g'_+ \left(\frac{x - y^+(x, t)}{t} \right) \geq g'_- \left(\frac{x - y^+(x, t)}{t} \right).$$

The other possibility is that $y^+(x + \varepsilon, t) > y^+(x, t)$ for all $\varepsilon > 0$. From right continuity of $x \mapsto y^+(x, t)$ it follows that $(x - y^+(x + \varepsilon, t))/t$ increases strictly to $(x - y^+(x, t))/t$ as $\varepsilon \searrow 0$. And then

$$\lim_{\varepsilon \searrow 0} g'_+ \left(\frac{x - y^+(x + \varepsilon, t)}{t} \right) = g'_- \left(\frac{x - y^+(x, t)}{t} \right).$$

In either case we obtain the second inequality of (3.4). \square

3.2. Differentiability of the flux and characteristics. In Corollary 3.1(b) below we show that differentiability of f implies $y^-(x, t) = y^+(x, t)$ for all but countably many x , the property used in part (3.2) of the Lax-Oleinik formula. In Proposition 3.1 we prove the property needed for Theorem 4(b), namely that $x^-(b; s, t) = x^+(b; s, t)$ for $0 < s < t$. Example 3.1 shows how various properties can fail without differentiability of f . The hypothesis of differentiability of f is used in the equivalent form that g is strictly convex on $[-1, 1]$.

Let $I(x; s, t)$ denote the set of $y \in [x - (t - s), x + (t - s)]$ at which the supremum in (2.15) is achieved.

Lemma 3.2. *Suppose $0 \leq t_1 < t_2 < t_3$, $y_2 \in I(x; t_2, t_3)$, and $y_1 \in I(y_2; t_1, t_2)$. Then $y_1 \in I(x; t_1, t_3)$. And either*

$$\frac{x - y_1}{t_3 - t_1} = \frac{x - y_2}{t_3 - t_2} = \frac{y_2 - y_1}{t_2 - t_1}$$

or g is linear between $(y_2 - y_1)/(t_2 - t_1)$ and $(x - y_2)/(t_3 - t_2)$.

Proof. Modify the proof of Rezakhanlou's (1995) Lemma 3.1. \square

Lemma 3.3. *Suppose g is strictly convex on $[-1, 1]$, $x_1 < x_2$, $y_i \in I(x_i; s, t)$ for $i = 1, 2$. Then $y_1 \leq y_2$.*

Proof. The assumptions $y_i \in I(x_i; s, t)$ give

$$u(y_1, s) - (t - s)g\left(\frac{x_1 - y_1}{t - s}\right) \geq u(y_2, s) - (t - s)g\left(\frac{x_1 - y_2}{t - s}\right)$$

and

$$u(y_2, s) - (t - s)g\left(\frac{x_2 - y_2}{t - s}\right) \geq u(y_1, s) - (t - s)g\left(\frac{x_2 - y_1}{t - s}\right).$$

To get a contradiction, suppose $y_1 - y_2 > 0$. Then the above inequalities give

$$\frac{g\left(\frac{x_2 - y_2}{t - s}\right) - g\left(\frac{x_2 - y_1}{t - s}\right)}{\left(\frac{x_2 - y_2}{t - s}\right) - \left(\frac{x_2 - y_1}{t - s}\right)} \leq \frac{g\left(\frac{x_1 - y_2}{t - s}\right) - g\left(\frac{x_1 - y_1}{t - s}\right)}{\left(\frac{x_1 - y_2}{t - s}\right) - \left(\frac{x_1 - y_1}{t - s}\right)}.$$

Since $x_2 - y_2 > x_1 - y_2$, $x_2 - y_1 > x_1 - y_1$, and $(x_i - y_i)/(t - s) \in [-1, 1]$, this contradicts the assumption on g . \square

Corollary 3.1. *Suppose g is strictly convex on $[-1, 1]$.*

- (a) *For $x_1 < x_2$, $y^+(x_1; s, t) \leq y^-(x_2; s, t)$.*
- (b) *For fixed $0 \leq s < t$, $y^-(x; s, t) = y^+(x; s, t)$ for all but countably many x .*
- (c) *$x^+(b; s, t) = \inf\{x : y^+(x; s, t) > b\}$ and $x^-(b; s, t) = \sup\{x : y^-(x; s, t) < b\}$.*
- (d) *If $x^-(b; s, t) < z < x^+(b; s, t)$, then $y^\pm(z; s, t) = b$.*

Proof. (a) is immediate from Lemma 3.3.

(b) Suppose x_0 is a continuity point of the nondecreasing, right-continuous function $x \mapsto y^+(x; s, t)$. Then

$$y^+(x_0; s, t) \geq y^-(x_0; s, t) \geq \lim_{x \nearrow x_0} y^+(x; s, t) = y^+(x_0; s, t),$$

where the second inequality follows from (a). Thus $y^-(x; s, t) = y^+(x; s, t)$ at all continuity points of $y^+(x; s, t)$, which includes all but countably many x .

(c) Let $w(b) = \inf\{x : y^+(x; s, t) > b\}$. We have $x^+(b; s, t) \geq w(b)$ without any new assumptions: if $x < w(b)$, then

$$y^+(x; s, t) \leq b \implies y^-(x; s, t) \leq b \implies x^+(b; s, t) \geq x.$$

For the converse we need part (a). Let $x > w(b)$. Then there exists x_1 such that $x > x_1 > w(b)$.

$$y^+(x_1; s, t) > b \implies y^-(x; s, t) > b \implies x^+(b; s, t) \leq x.$$

We leave the proof of the other formula to the reader.

(d) Immediate from definition (2.18–19) and part (c). \square

Proposition 3.1. *Suppose g is strictly convex on $[-1, 1]$, and $s > 0$. Then $x^-(b; s, t) = x^+(b; s, t)$ for all $b \in \mathbf{R}$ and $t > s$.*

Proof. Suppose there exists a z such that $x^-(b; s, t) < z < x^+(b; s, t)$. By Corollary (d) $y^\pm(z; s, t) = b$, and consequently for $\varepsilon > 0$,

$$u(b, s) - (t - s)g\left(\frac{z - b}{t - s}\right) \geq u(b \pm \varepsilon, s) - (t - s)g\left(\frac{z - b \mp \varepsilon}{t - s}\right).$$

Letting $\varepsilon \searrow 0$ then gives

$$\limsup_{\varepsilon \searrow 0} \frac{u(b + \varepsilon, s) - u(b, s)}{\varepsilon} \leq -g'_-\left(\frac{z - b}{t - s}\right)$$

and

$$\liminf_{\varepsilon \searrow 0} \frac{u(b - \varepsilon, s) - u(b, s)}{-\varepsilon} \geq -g'_+\left(\frac{z - b}{t - s}\right).$$

Since we may vary z in an interval, and since by assumption the slope of g cannot remain constant in this interval, we conclude the strict inequality

$$\limsup \frac{u(b + \varepsilon, s) - u(b, s)}{\varepsilon} < \liminf \frac{u(b - \varepsilon, s) - u(b, s)}{-\varepsilon}.$$

By (3.3–4), this implies

$$(3.5) \quad g'_+ \left(\frac{b - y^+(b, s)}{s} \right) > g'_- \left(\frac{b - y^-(b, s)}{s} \right).$$

The slope of a convex function is nondecreasing and by definition $(b - y^-(b, s))/s \geq (b - y^+(b, s))/s$. So (3.5) implies that $y^-(b, s) = y^+(b, s)$ and g has a corner at $(b - y^\pm(b, s))/s$. Since we are assuming g strictly convex in $[-1, 1]$, Lemma 3.2 now forces $(z - b)/(t - s) = (b - y^\pm(b, s))/s$. This cannot hold for distinct z 's. But the preceding argument is valid for all $z \in (x^-(b; s, t), x^+(b; s, t))$, so to avoid contradiction we must have $x^-(b; s, t) = x^+(b; s, t)$. \square

Finally we show by example how a point of nondifferentiability in f destroys all the good properties of the characteristics.

Example 3.1. Suppose $\rho \in (0, K)$ is a point such that $f'_-(\rho) = \xi_2 > \xi_1 = f'_+(\rho)$. Then g has a linear segment of slope $-\rho$ over $[\xi_1, \xi_2]$. Apply the formulas to the constant initial profile $\rho_0(x) \equiv \rho$, with $u_0(x) = \rho x$. We get $I(x; s, t) = [x - \xi_2(t - s), x - \xi_1(t - s)]$, $y^-(x; s, t) = x - \xi_2(t - s)$, $y^+(x; s, t) = x - \xi_1(t - s)$, $x^-(b; s, t) = b + \xi_1(t - s)$, $x^+(b; s, t) = b + \xi_2(t - s)$, and $u(x, t) = \rho x - tf(\rho)$. In particular $\rho(x, t) = u_x(x, t) \equiv \rho$, so the constant profile is the solution. But the conclusions of Lemma 3.3, Corollary 3.1(a)–(b), and Proposition 3.1 have been contradicted.

3.3. Proof of Theorem 2. We prove the uniqueness criterion of Theorem 2 under these general assumptions: The flux f is continuous and concave on an interval I , and g is defined by $g(x) = \sup_{\rho \in I} \{f(\rho) - \rho x\}$. The initial density $\rho_0(x)$ is a given locally bounded measurable function on \mathbf{R} , $u(x, t)$ is a locally Lipschitz function on $\mathbf{R} \times [0, \infty)$ defined by the Hopf-Lax formula (2.10), and $\rho(x, t) = u_x(x, t)$ exists for a.e. x for any fixed t . $\lambda(x, t)$ is a locally bounded measurable function on $\mathbf{R} \times [0, \infty)$ that satisfies the boundary condition $\lambda(x, 0) = \rho_0(x)$ a.e. and the integral condition (2.13). The ranges of ρ and λ lie in I .

The goal is to show that, for a.e. x ,

$$(3.6) \quad \int_0^t f(\lambda(x, s)) ds \leq \int_0^t f(\rho(x, s)) ds$$

for all $t \geq 0$. And conversely, if equality holds in (3.6) for almost all (x, t) , then $\lambda(x, t) = \rho(x, t)$ for almost all (x, t) .

Some technical difficulties stem from assuming only (2.13) and no regularity on λ . We cannot expect the function $x \mapsto \lambda(x, t)$ to be a sensible object for every fixed t because $\lambda(\cdot, t)$ can be redefined on any null set of t 's without affecting (2.13). At $t = 0$ we have

$$(3.7) \quad \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_0^\varepsilon dt \int_{\mathbf{R}} \lambda(x, t) \psi(x) dx = \int_{\mathbf{R}} \rho_0(x) \psi(x) dx$$

for $\psi \in C_c^1(\mathbf{R})$. This is obtained from (2.13) by taking $\phi(x, t) = \psi(x)g(t)$ where $g(0) = 1$, $g' = -1/\varepsilon$ on $(0, \varepsilon)$, and $g(t) = 0$ for $t \geq \varepsilon$. [The C^1 test functions in (2.13) can be replaced by continuous, piecewise C^1 test functions by taking limits]

A corresponding property holds for certain later times t . Abbreviate

$\text{Leb}(h)$ = the set of Lebesgue points of the function h

for any measurable, locally bounded function h on either \mathbf{R} or $\mathbf{R} \times [0, \infty)$. [For the definition and properties of Lebesgue points see Section 7.2 in Wheeden and Zygmund (1977).] Define

$$\mathbf{T} = \{t > 0 : (x, t) \in \text{Leb}(\lambda) \text{ for almost every } x \in \mathbf{R}\}.$$

Since the complement of $\text{Leb}(\lambda)$ is a null set of $\mathbf{R} \times (0, \infty)$, almost every t lies in \mathbf{T} . By considering test functions of the type $\phi(x, t) = \psi(x)g(t)$ for suitable g , one obtains from (2.13) a limit such as (3.7) around $t \in \mathbf{T}$, and then the following for $t \in \mathbf{T}$:

$$(3.8) \quad \int_{\mathbf{R}} \psi(x) \lambda(x, t) dx - \int_{\mathbf{R}} \psi(x) \rho_0(x) dx = \int_0^t \int_{\mathbf{R}} f(\lambda(x, s)) \psi'(x) dx ds$$

for compactly supported C^1 functions ψ on \mathbf{R} . By taking limits of such functions, (3.8) is valid for compactly supported, continuous, piecewise C^1 functions ψ .

Lemma 3.4. (i) *Fix finite $A < B$ and $T > 0$. Then there exists a constant $C = C(A, B, T)$ such that, for all $A \leq a < b \leq B$ and $s, t \in \mathbf{T} \cup \{0\}$ such that $0 \leq s, t \leq T$,*

$$(3.9) \quad \left| \int_a^b \lambda(x, t) dx - \int_a^b \lambda(x, s) dx \right| \leq C \cdot |t - s|.$$

(ii) *If $t \in \mathbf{T} \cup \{0\}$ and $(a, t), (b, t) \in \text{Leb}(f \circ \lambda)$, then we have a time derivative along points of \mathbf{T} :*

$$(3.10) \quad \lim_{\substack{s \rightarrow t \\ s \in \mathbf{T}}} \frac{1}{t - s} \left\{ \int_a^b \lambda(x, t) dx - \int_a^b \lambda(x, s) dx \right\} = f(\lambda(a, t)) - f(\lambda(b, t)).$$

Proof. In (3.8) take ψ such that $\psi = 0$ outside $[a - \varepsilon, b + \varepsilon]$, $\psi' = 1/\varepsilon$ on $(a - \varepsilon, a)$, $\psi = 1$ on $[a, b]$, and $\psi' = -1/\varepsilon$ on $(b, b + \varepsilon)$. We get

$$(3.11) \quad \begin{aligned} & \int_a^b \lambda(x, t) dx - \int_a^b \lambda(x, s) dx + O(\varepsilon) \\ &= \frac{1}{\varepsilon} \int_s^t \int_{a-\varepsilon}^a f(\lambda(x, \tau)) dx d\tau - \frac{1}{\varepsilon} \int_s^t \int_b^{b+\varepsilon} f(\lambda(x, \tau)) dx d\tau. \end{aligned}$$

The $O(\varepsilon)$ term contains the integrals over $[a - \varepsilon, a]$ and $[b, b + \varepsilon]$ from the left-hand side. These are bounded by $O(\varepsilon)$ uniformly over the range of a, b, s, t by the local boundedness of λ . The $1/\varepsilon$ factors on the right are the slopes of ψ . The second line in (3.11) is bounded by $C|t - s|$ in absolute value, where C is twice the supremum of $|f(\lambda(x, \tau))|$ over $[A, B] \times [0, T]$. Part (i) follows by letting $\varepsilon \searrow 0$. For part (ii) set $\varepsilon = \delta|t - s|$, divide through (3.11) by $t - s$, let first $\mathbf{T} \ni s \rightarrow t$ and then $\delta \searrow 0$. \square

As a locally Lipschitz function $u(x, t)$ is differentiable a.e. by Rademacher's theorem [Section 5.8.3 in Evans (1998)]. And as the viscosity solution defined by the Hopf-Lax formula, $u(x, t)$ satisfies the equation $u_t + f(u_x) = 0$ a.e. [Theorem 5, Section 3.3.2 in Evans (1998)]. By definition $\rho = u_x$ exists a.e., so for almost every x_0 and all $t > 0$,

$$(3.12) \quad \int_0^t f(\rho(x_0, s)) ds = - \int_0^t u_t(x_0, s) ds = u_0(x_0) - u(x_0, t).$$

It is also the case that for almost every x_0 , $(x_0, t) \in \text{Leb}(f \circ \lambda)$ for almost every t . Fix such an x_0 for which also (3.12) holds. For $(x, t) \in \mathbf{R} \times \mathbf{T}$ define

$$(3.13) \quad v(x, t) = \int_{x_0}^x \lambda(y, t) dy - \int_0^t f(\lambda(x_0, s)) ds + u_0(x_0).$$

The first integral is interpreted with a sign so that $v(x_2, t) - v(x_1, t) = \int_{x_1}^{x_2} \lambda(y, t) dy$ for $x_1 < x_2$. By Lemma 3.4(i) and local boundedness of λ and $f \circ \lambda$, v is locally Lipschitz on $\mathbf{R} \times \mathbf{T}$. Consequently v extends uniquely to a locally Lipschitz function on $\mathbf{R} \times [0, \infty)$. This extension has the correct boundary values at $t = 0$ because by Lemma 3.4(i),

$$\begin{aligned} \lim_{\mathbf{T} \ni t \rightarrow 0} v(x, t) &= \int_{x_0}^x \lambda(y, 0) dy + u_0(x_0) = \int_{x_0}^x \rho_0(y) dy + u_0(x_0) \\ &= u_0(x). \end{aligned}$$

The goal is now to prove

$$(3.14) \quad v(x, t) \geq u(x, t) \text{ for all } (x, t).$$

Setting $x = x_0$ in (3.14) then gives, by (3.12) and (3.13), the conclusion (3.6) for $x = x_0$ and all $t \in \mathbf{T}$. Both sides of (3.6) are continuous in t because the integrands are locally bounded, so (3.6) follows for all t . This whole argument beginning with (3.12) can be repeated for almost every x_0 . At the end of this section we argue that equality in (3.6) implies $\lambda = \rho$ a.e. First we prove (3.14).

Fix (x, t) . For a fixed $y \in \mathbf{R}$ set $\xi(s) = y + (s/t)(x - y)$ for $s \in [0, t]$. As y varies we consider the collection of line segments $\{(\xi(s), s) : 0 \leq s \leq t\}$ on the plane $\mathbf{R} \times [0, \infty)$. Their union forms the “backward light cone” from the apex (x, t) to the base line $\mathbf{R} \times \{0\}$. For almost every such y , it must be that

$$(\xi(s), s) \in \text{Leb}(f \circ \lambda) \cap \text{Leb}(\lambda)$$

for almost every $s \in [0, t]$, because only a null set of points of $\mathbf{R} \times [0, \infty)$ fail to be Lebesgue points. Fix now such a y .

The function $\gamma(s) = v(\xi(s), s)$, $0 \leq s \leq t$, is Lipschitz, and hence $\gamma'(s)$ exists for a.e. s and

$$(3.15) \quad \gamma(t) - \gamma(0) = \int_0^t \gamma'(s) ds.$$

Fix $s_0 \in [0, t]$ such that all this holds: $s_0 \in \mathbf{T}$, $\gamma'(s_0)$ exists, $(x_0, s_0), (\xi(s_0), s_0) \in \text{Leb}(f \circ \lambda) \cap \text{Leb}(\lambda)$, and $s_0 \in \text{Leb}\{f(\lambda(x_0, \cdot))\}$. These conditions are satisfied by a.e. s_0 . For such an s_0 we prove that

$$(3.16) \quad \gamma'(s_0) = \frac{x-y}{t} \lambda(\xi(s_0), s_0) - f(\lambda(\xi(s_0), s_0)).$$

To prove (3.16) we calculate $\gamma'(s_0)$ as the limit

$$\lim_{\substack{s \searrow s_0 \\ s \in \mathbf{T}}} \frac{\gamma(s) - \gamma(s_0)}{s - s_0}.$$

So consider $s \in \mathbf{T}$, $s > s_0$.

$$\begin{aligned} \frac{\gamma(s) - \gamma(s_0)}{s - s_0} &= \frac{v(\xi(s), s) - v(\xi(s_0), s_0)}{s - s_0} \\ &= \frac{1}{s - s_0} \int_{\xi(s_0)}^{\xi(s)} \lambda(y, s) dy + \frac{1}{s - s_0} \left\{ \int_{x_0}^{\xi(s_0)} \lambda(y, s) dy - \int_{x_0}^{\xi(s_0)} \lambda(y, s_0) dy \right\} \\ &\quad - \frac{1}{s - s_0} \int_{s_0}^s f(\lambda(x_0, \tau)) d\tau \\ &\equiv A_1 + A_2 - A_3. \end{aligned}$$

Term by term: For A_1 first use $s - s_0 = (\xi(s) - \xi(s_0))t/(x - y)$. By the assumption $s \in \mathbf{T}$ and Lemma 3.4(i), we can replace s by an average over $[s_0, s] \cap \mathbf{T}$ at the expense of an $O(s - s_0)$ error term. Since \mathbf{T} covers almost all of $[s_0, s]$ we can ignore the restriction to \mathbf{T} in the average. These steps give

$$\begin{aligned} A_1 &= \frac{1}{s - s_0} \int_{\xi(s_0)}^{\xi(s)} \lambda(y, s) dy \\ &= \frac{x-y}{t} \cdot \frac{1}{(s - s_0)(\xi(s) - \xi(s_0))} \int_{s_0}^s d\tau \int_{\xi(s_0)}^{\xi(s)} \lambda(y, \tau) dy + O(s - s_0) \\ &\longrightarrow \frac{x-y}{t} \lambda(\xi(s_0), s_0) \end{aligned}$$

as $s \searrow s_0$ along points of \mathbf{T} , by the assumption $(\xi(s_0), s_0) \in \text{Leb}(\lambda)$. Next,

$$A_2 \longrightarrow f(\lambda(x_0, s_0)) - f(\lambda(\xi(s_0), s_0))$$

as $s \searrow s_0$ along points of \mathbf{T} , by the assumptions $s_0 \in \mathbf{T}$ and $(x_0, s_0), (\xi(s_0), s_0) \in \text{Leb}(f \circ \lambda)$ and by Lemma 3.4(ii). Finally,

$$A_3 \longrightarrow f(\lambda(x_0, s_0))$$

by the assumption $s_0 \in \text{Leb}\{f(\lambda(x_0, \cdot))\}$. Altogether we have proved (3.16) for a.e. s_0 .

By (2.8), (3.16) implies $\gamma'(s) \geq -g((x - y)/t)$ for a.e. s . Since $v(y, 0) = u_0(y)$, we can rewrite (3.15) as

$$v(x, t) = u_0(y) + \int_0^t \gamma'(s) ds \geq u_0(y) - tq((x - y)/t).$$

This argument is valid for a.e. y , and consequently by the Hopf-Lax formula (2.10), $v(x, t) \geq u(x, t)$. We have proved (3.14), and thereby (3.6) for $x = x_0$. The choice of x_0 was such that it covers a.e. point in \mathbf{R} .

For the converse, suppose

$$(3.17) \quad \int_0^t f(\lambda(x, s))ds = \int_0^t f(\rho(x, s))ds \quad \text{for a.e. } (x, t).$$

(3.8) is valid for both λ and ρ for a.e. t . Consider such a t for which also (3.17) holds for a.e. x . The right-hand side of (3.8) is $\int_{\mathbf{R}} dx \psi(x) \int_0^t f(\lambda(x, s))ds$, so by (3.17) its value is not changed by replacing λ with ρ . We conclude that for a.e. t , $\int_{\mathbf{R}} \lambda(x, t)\psi(x)dx = \int_{\mathbf{R}} \rho(x, t)\psi(x)dx$ for all test functions ψ . This implies $\lambda = \rho$ a.e.

4. SECOND CLASS PARTICLES IN THE VARIATIONAL COUPLING

4.1. Construction and variational coupling. First we construct the process. Let $\{D_i : i \in \mathbf{Z}\}$ be a collection of mutually independent rate 1 Poisson jump time processes on the time line $(0, \infty)$, defined on some probability space (Ω, \mathcal{F}, P) . We construct the process $\eta(t) = (\eta(i, t) : i \in \mathbf{Z})$ on Ω in terms of “current particles.” These form a process $z(t) = (z(i, t) : i \in \mathbf{Z})$ of labeled particles that move on \mathbf{Z} . The location of the i th particle at time t is $z(i, t)$, and these satisfy

$$(4.1) \quad 0 \leq z(i+1, t) - z(i, t) \leq K.$$

In the graphical construction, $z(i)$ attempts to jump 1 step to the *left* at epochs of D_i . If the jump violates (4.1), it is suppressed. We can summarize the jump rule like this:

$$(4.2) \quad \begin{aligned} &\text{If } t \text{ is an epoch of } D_i, \text{ then} \\ &z(i, t) = \max\{z(i, t-) - 1, z(i-1, t-), z(i+1, t-) - K\}. \end{aligned}$$

To construct $z(\cdot)$ from rule (4.2) we exclude an exceptional null set of realizations of $\{D_i\}$. For the remainder of this section, fix a realization $\{D_i\}$ with these properties:

- (4.3a) There are no simultaneous jump attempts.
- (4.3b) Each D_i has only finitely many epochs in every bounded time interval.
- (4.3c) Given any $t_1 > 0$, there are arbitrarily faraway indices $i_0 \ll 0 \ll i_1$ such that D_{i_0} and D_{i_1} have no epochs in the time interval $[0, t_1]$.

These properties are satisfied almost surely, so the evolution $z(t)$, $0 \leq t < \infty$, is well-defined almost everywhere on Ω . Then the process $\eta(t)$ is defined by

$$(4.4) \quad \eta(i, t) = z(i, t) - z(i-1, t).$$

It should be clear that $\eta(\cdot)$ satisfies the rules described in Section 2.

Define a family $\{w^k : k \in \mathbf{Z}\}$ of auxiliary processes on Ω . Each $w^k(t) = (w^k(i, t) : i \in \mathbf{Z})$ is a process of the same type as $z(t)$, so (4.1) is in force for all $w^k(i, t)$. The initial configuration of w^k depends on the initial position $z(k, 0)$:

$$w^k(i, 0) = \begin{cases} z(k, 0), & i \geq 0 \\ (1, \infty) + K, & i < 0 \end{cases}$$

The process w^k reads its jump commands from the same Poisson processes $\{D_i\}$, but with a translated index: at epochs t of D_{i+k} ,

$$w^k(i, t) = \max\{w^k(i, t-) - 1, w^k(i-1, t-), w^k(i+1, t-) - K\}.$$

We prove a number of pathwise properties for the processes z and $\{w^k\}$ that are valid under the fixed realization $\{D_i\}$ that satisfies (4.3a–c). Since (4.3a–c) are valid almost surely, the properties we derive are almost sure properties, but we leave the modifier “almost surely” out of the statements that follow.

The coupling of the processes z and $\{w^k\}$ preserves ordering. One consequence is the following inequality, valid for all $k < l$, all i and all t :

$$(4.5) \quad w^l(i-l, t) \leq w^k(i-k, t) + [z(l, 0) - z(k, 0)].$$

The usefulness of the family of processes $\{w^k\}$ lies in this “variational coupling” lemma:

Lemma 4.1. *For all $i \in \mathbf{Z}$ and $t \geq 0$,*

$$(4.6) \quad z(i, t) = \sup_{k \in \mathbf{Z}} w^k(i-k, t).$$

In the case $K = \infty$ the supremum in (4.6) is over $\{k : k \leq i\}$ because $w^k(i-k, t) = -\infty$ for $k > i$.

For some purposes it is convenient to modify the processes w^k so that they start at level zero and advance in the positive direction. So define processes $\{\xi^k\}$ by

$$\xi^k(i, t) = z(k, 0) - w^k(i, t).$$

The process ξ^k does not depend on $z(k, 0)$, and depends on the superscript k only through a translation of the i -index of the Poisson processes $\{D_i\}$. Initially

$$\xi^k(i, 0) = \begin{cases} 0, & i \geq 0 \\ -Ki, & i < 0. \end{cases}$$

Dynamically, at epochs t of D_{i+k} ,

$$\xi^k(i, t) = \min\{\xi^k(i, t-) + 1, \xi^k(i-1, t-), \xi^k(i+1, t-) + K\}.$$

These jumps preserve the inequalities

$$\xi^k(i, t) \leq \xi^k(i-1, t) \quad \text{and} \quad \xi^k(i, t) \leq \xi^k(i+1, t) + K.$$

Now (4.6) can be written as

$$(4.7) \quad z(i, t) = \sup_{k \in \mathbf{Z}} \{z(k, 0) - \xi^k(i-k, t)\}.$$

This variational formula is a microscopic analogue of the Hopf-Lax formula (2.10), and the basis for the proof of the hydrodynamic limit. To prove Theorem 1, one shows that the right-hand side of (4.7) converges to the right-hand side of (2.10). For more details about the construction, the variational coupling lemma, and the proofs, the reader may consult Sennäläinen (1999).

4.2. The second class particle. We investigate the position $X(t)$ of a second class particle in the variational coupling. Fix again a sample point so that we can regard $\{D_i\}$, $(\eta(i, 0))$, and $X(0)$ as deterministic, with (4.3a-c) satisfied by $\{D_i\}$.

As defined in Section 2.2, $X(t)$ is the location of the unique discrepancy between two processes η and $\tilde{\eta}$ that satisfy initially $\tilde{\eta}(X(0), 0) = \eta(X(0), 0) + 1$ and $\tilde{\eta}(i, 0) = \eta(i, 0)$ for $i \neq X(0)$. In particular, the second class particle is not counted in the η -variables and by the K -exclusion rule always $\eta(X(t), t) \leq K - 1$.

We define η and $\tilde{\eta}$ by (4.4), in terms of processes z and \tilde{z} that initially satisfy

$$\tilde{z}(i, 0) = z(i, 0) \text{ for } i \leq X(0) - 1 \text{ and } \tilde{z}(i, 0) = z(i, 0) + 1 \text{ for } i \geq X(0).$$

Processes z and \tilde{z} follow the same Poisson processes $\{D_i\}$, so this is the basic coupling. The location $X(t)$ is always uniquely defined as the location of the discrepancy, as shown in this lemma:

Lemma 4.2. *For all $t \geq 0$ there is a finite location $X(t)$ such that*

$$(4.8) \quad \tilde{z}(i, t) = z(i, t) \text{ for } i \leq X(t) - 1 \text{ and } \tilde{z}(i, t) = z(i, t) + 1 \text{ for } i \geq X(t).$$

Proof. Given any finite time horizon t_0 , find indices $\dots < i_{-1} < i_0 < i_1 < \dots$ such that D_{i_m} has no jump epochs in the time interval $(0, t_0]$ for all m , $i_m \rightarrow \pm\infty$ as $m \rightarrow \pm\infty$, and $i_0 < X(0) < i_1$. Particles $z(i_m)$ and $\tilde{z}(i_m)$ do not move up to time t_0 . In each finite segment (i_{m-1}, i_m) there are only finitely many jump epochs up to time t_0 . Arrange these finitely many jump epochs in temporal order, and then prove by induction that the lemma holds after each jump time. \square

(4.8) defines $X(t)$ in our framework. Next we give $X(t)$ a representation that is a microscopic analogue of the characteristics.

Proposition 4.1. *We have the formulas*

$$(4.9) \quad X(t) = \inf\{i \in \mathbf{Z} : z(i, t) = z(k, 0) - \xi^k(i - k, t) \text{ for some } k \geq X(0)\}$$

and

$$(4.10) \quad X(t) = 1 + \sup\{i \in \mathbf{Z} : \tilde{z}(i, t) = \tilde{z}(k, 0) - \xi^k(i - k, t) \text{ for some } k < X(0)\}.$$

Proof. Consider these four statements:

$$(4.11) \quad \text{If } i < X(t), \text{ then } z(i, t) > z(k, 0) - \xi^k(i - k, t) \text{ for all } k \geq X(0).$$

$$(4.12) \quad \text{If } i \geq X(t), \text{ then } z(i, t) = z(k, 0) - \xi^k(i - k, t) \text{ for some } k \geq X(0).$$

$$(4.13) \quad \text{If } i < X(t), \text{ then } \tilde{z}(i, t) = \tilde{z}(k, 0) - \xi^k(i - k, t) \text{ for some } k < X(0).$$

$$(4.14) \quad \text{If } i \geq X(t), \text{ then } \tilde{z}(i, t) > \tilde{z}(k, 0) - \xi^k(i - k, t) \text{ for all } k < X(0).$$

These statements imply that the infimum and supremum in the formulas (4.9–10) are well-defined and finite, and also prove the formulas themselves.

To contradict (4.11), suppose $i < X(t)$ and

$$(4.15) \quad z(i, t) = z(k, 0) - \xi^k(i - k, t)$$

for some $k \geq X(0)$. Then by (4.8)

$$\tilde{z}(i, t) = \tilde{z}(k, 0) - \xi^k(i - k, t) - 1$$

which contradicts the variational formula (4.7) for process \tilde{z} . This contradiction proves (4.11), and also implies that there must exist some $k < X(0)$ such that (4.15) holds. By (4.8), (4.15) then turns into

$$\tilde{z}(i, t) = \tilde{z}(k, 0) - \xi^k(i - k, t),$$

and this proves (4.13).

(4.12) and (4.14) are proved by a similar argument that starts by contradicting (4.14). \square

The second class particle may jump either left or right, but a certain monotonicity can be found from the representation (4.9):

Proposition 4.2. *Let*

$$k_1(t) = \sup\{k \geq X(0) : z(X(t), t) = w^k(X(t) - k, t)\}$$

denote the maximal k that satisfies the requirement in formula (4.9), or infinity. Then $k_1(t)$ is nondecreasing as a function of time. If there are initially arbitrarily large indices j such that $\eta(j, 0) \leq K - 1$, then $k_1(t)$ is finite for all t .

We shall not prove Proposition 4.2 for we make no use of it in this paper.

5. PROOFS OF THE LAWS OF LARGE NUMBERS

Now assume we are in the setting described in the paragraph preceding Theorem 1. The initial particle configurations $(\eta_n(i, 0) : i \in \mathbf{Z})$, $n \in \mathbf{N}$, and the Poisson jump time processes $\{D_i\}$ are defined on a probability space (Ω, \mathcal{F}, P) . Define initial configurations $(z_n(i, 0) : i \in \mathbf{Z})$ by setting $z_n(0, 0) = 0$ and using (4.4). Construct the processes $z_n(\cdot)$ as in Section 4.1, and then define the processes $\eta_n(\cdot)$ by (4.4) in terms of $z_n(\cdot)$. The location of the second class particle is given by the variational representation in Proposition 4.1: for the n th process

$$\begin{aligned} X_n(t) &= \inf\{i \in \mathbf{Z} : z_n(i, t) = w_n^k(i - k, t) \text{ for some } k \geq X_n(0)\} \\ &= \inf\{i \in \mathbf{Z} : z_n(i, t) = z_n(k, 0) - \xi^k(i - k, t) \\ &\quad \text{for some } k \geq X_n(0)\}. \end{aligned}$$

Theorems 3 and 4 rely on the basic hydrodynamic limits, which we summarize here from Seppäläinen (1999) in terms of the z -variables. The following statements all hold almost surely:

$$(5.1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} z_n([ny], 0) = u_0(u) \quad \text{for all } u \in \mathbf{R}:$$

$$(5.2) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \xi^{[nr]}([nx], nt) = tg(x/t) \quad \text{for all } x, r \in \mathbf{R} \text{ and } t > 0;$$

and

$$(5.3) \quad \lim_{n \rightarrow \infty} \frac{1}{n} z_n([nx], nt) = u(x, t) \quad \text{for all } x \in \mathbf{R} \text{ and } t > 0.$$

The function u_0 is defined by (2.7) from the given initial profile ρ_0 . (5.1) is a restatement of assumption (2.3). (5.2) is proved by subadditivity, and this limit defines g . Then $u(x, t)$ is defined in terms of u_0 and g by the Hopf-Lax formula (2.10). Next (5.3) is proved, by (5.1), (5.2), and the variational coupling (4.7). Lastly, f is defined as the conjugate of g , and $u(x, t)$ is identified as the viscosity solution of (2.11).

The following estimate [Seppäläinen (1999), Lemma 6.1] reduces the range of indices needed in the variational formula. It will be used several times in the proofs.

Lemma 5.1. *Define*

$$\zeta_n(i, l, t) = \max_{i-l \leq k \leq i+l} \{z_n(k, 0) - \xi^k(i - k, t)\}.$$

Fix $r > t$. Then there exists a finite positive constant $C = C(r, t)$ such that for all $i \in \mathbf{Z}$ and all $n \geq 1$,

$$(5.4) \quad P(z_n(i, nt) \neq \zeta_n(i, nr, nt)) \leq e^{-Cn}.$$

First we prove Theorem 3, so recall assumption (2.20):

$$(5.5) \quad \lim_{n \rightarrow \infty} \frac{1}{n} X_n(0) = b \quad \text{a.s.}$$

Proposition 5.1. *Assume (2.3) and (2.20). Then*

$$\limsup_{n \rightarrow \infty} n^{-1} X_n(nt) \leq x^+(b, t) \quad \text{almost surely.}$$

Proof. Fix $x_1 > x^+(b, t)$. We shall show that $\limsup_{n \rightarrow \infty} n^{-1} X_n(nt) \leq x_1$ almost surely. By definition $x^+(b, t) = \sup\{x : y^-(x, t) \leq b\}$, so $y^-(x_1, t) > b$. Choose c_1 so that $b < c_1 < y^-(x_1, t)$.

Let Γ be the event on which

$$(5.6) \quad X_n(nt) > nx_1 \text{ for infinitely many } n,$$

assumptions (4.3a–c) of Section 4.1 hold so that we can use variational formulas (4.7) and (4.9), the laws of large numbers (5.1)–(5.3) and (5.5) hold, and

$$(5.7) \quad z_n([nx_1], nt) = w_n^k([nx_1] - k, nt)$$

for some $k \geq [nc_0]$ for all large enough n , for some fixed finite c_0 . Except for (5.6) the conditions of Γ hold with probability one, so it suffices to show that Γ is empty.

Suppose Γ is not empty. Fix a sample point in Γ , and suppose n is such that $X_n(nt) > nx_1$. Then $z_n([nx_1], nt) \neq w_n^k([nx_1] - k, nt)$ for all $k \geq X_n(0)$, so there must exist a $k < X_n(0)$ such that (5.7) holds. Since limit (5.5) holds on Γ , $X_n(0) \leq [nc_1]$ for large enough n . Thus for a subsequence of n 's there exists a k in (5.7) such that $[nc_0] \leq k \leq [nc_1]$.

Pick a partition $c_0 = y_0 < y_1 < \dots < y_m = c_1$ of $[c_0, c_1]$ with mesh $\Delta y = \max(y_{i+1} - y_i)$. Fix i so that for infinitely many n in the subsequence picked above, some k in (5.7) satisfies $[ny_i] \leq k \leq [ny_{i+1}]$. By (4.5) we have, for infinitely many n ,

$$(5.8) \quad \begin{aligned} z_n([nx_1], nt) &= w_n^k([nx_1] - k, nt) \\ &\leq w_n^{[ny_i]}([nx_1] - [ny_i], nt) + \{z_n([ny_{i+1}], 0) - z_n([ny_i], 0)\}. \end{aligned}$$

Divide by n and let $n \rightarrow \infty$ along the appropriate subsequence of n 's. Since the laws of large numbers hold for our fixed sample point, we get

$$(5.9) \quad u(x_1, t) \leq u_0(y_i) - tg\left(\frac{x_1 - y_i}{t}\right) + \{u_0(y_{i+1}) - u_0(y_i)\}.$$

Take the mesh $\Delta y \rightarrow 0$. By continuity and compactness, we conclude that the Hopf-Lax formula (2.10) for $u(x_1, t)$ has a maximizer in $[c_0, c_1]$, so that $y^-(x_1, t) \leq c_1$. This contradiction with the choice of c_1 implies that Γ is empty. \square

Proposition 5.2. *Assume (2.3) and (2.20). Then*

$$\liminf_{n \rightarrow \infty} n^{-1} X_n(nt) \geq x^-(b, t) \quad \text{almost surely.}$$

Proof. Fix $x_1 < x^-(b, t)$. Consider the event Γ on which $X_n(nt) < nx_1$ for infinitely many n , assumptions (4.3a-c) hold, and the laws of large numbers (5.1)–(5.3), and (5.5) hold. We assume that Γ has positive probability, and derive a contradiction.

(4.12) and $X_n(nt) < nx_1$ imply that for some $k \geq X_n(0)$,

$$(5.10) \quad z_n([nx_1], nt) = z_n(k, 0) - \xi^k([nx_1] - k, nt).$$

As in Section 4.2, we have the processes \tilde{z}_n , coupled with z_n through the Poisson processes $\{D_i\}$, that satisfy

$$(5.11) \quad \begin{aligned} \tilde{z}_n(i, nt) &= z_n(i, nt) \text{ for } i < X_n(nt) \\ \text{and } \tilde{z}_n(i, nt) &= z_n(i, nt) + 1 \text{ for } i \geq X_n(nt). \end{aligned}$$

(4.10) implies that

$$(5.12) \quad \tilde{z}_n([nx_1], nt) = \tilde{z}_n(k, 0) - \xi^k([nx_1] - k, nt).$$

By (4.14), (5.12) cannot hold for any $k < X_n(0)$. By applying Lemma 5.1 to the processes \tilde{z}_n and by shrinking Γ by no more than a null set, we may assume that for large enough n the k in (5.12) must lie in the range $X_n(0) \leq k \leq [nc_1]$ for a constant c . Then by (5.5) we conclude that for large n , $[nc_0] \leq k \leq [nc_1]$ where c

can be taken arbitrarily close to b . By left-continuity of $x^-(\cdot, t)$ and the assumption $x_1 < x^-(b, t)$, we may choose c_0 so that $x_1 < x^-(c_0, t)$.

Now starting from (5.12) repeat the argument around (5.8)–(5.9) for \tilde{z}_n . By (5.11) \tilde{z}_n satisfies the limits (5.1) and (5.3). As in the step after (5.9), we get that $u(x_1, t)$ has a maximizer in $[c_0, c_1]$ so that $y^+(x_1, t) \geq c_0$. By definition (2.18) this implies $x^-(c_0, t) \leq x_1$ and contradicts the choice of c_0 . \square

Propositions 5.1–5.2 prove Theorem 3.

Proof of Theorem 4(a). Assume $x(y, t) = x^\pm(y, t)$ for all $y \in \mathbf{R}$. Let Γ be the event on which (4.3a–c) and (5.1)–(5.3) hold, and also for large enough n

$$(5.13) \quad -nb_0 \leq X_n(0) \leq nb_0$$

for some constant b_0 . We prove that

$$(5.14) \quad \lim_{n \rightarrow \infty} \left| \frac{X_n(nt)}{n} - x\left(\frac{X_n(0)}{n}, t\right) \right| = 0.$$

almost surely on Γ . By assumption (2.23), Theorem 4(a) follows by repeating this for a sequence of b_0 's increasing to ∞ .

Since $x^-(y, t)$ is left-continuous and $x^+(y, t)$ right-continuous in y , the assumed unique characteristic $x(y, t) = x^\pm(y, t)$ is continuous in y . Given $\varepsilon > 0$, choose a partition $-b_0 = r_0 < r_1 < \dots < r_h = b_0$ so that $x(r_{j+1}, t) < x(r_j, t) + \varepsilon/2$. Set $x_j = x(r_j, t) + \varepsilon$. Shrink Γ by a null set to guarantee that a.s. for every $j = 0, \dots, h$, for large enough n there exists a $k \geq nc_0$ such that

$$z_n([nx_j], nt) = w_n^k([nx_j] - k, nt).$$

This can be done by Lemma 5.1 if c_0 is chosen small enough.

To contradict (5.14), suppose Γ contains a sample point for which

$$(5.15) \quad X_n(nt) > nx(n^{-1}X_n(0), t) + n\varepsilon$$

for infinitely many n . Restrict this subsequence of n 's further so that for some fixed j ,

$$(5.16) \quad nr_j \leq X_n(0) \leq nr_{j+1}$$

for all these n . By the monotonicity of $x(\cdot, t)$, (5.15–16) imply that $X_n(nt) > nx_j$ for infinitely many n . For each such n there exists $k < X_n(0)$ and no $k \geq X_n(0)$ such that $z_n([nx_j], nt) = w_n^k([nx_j] - k, nt)$. By the assumptions made above, $nc_0 \leq k \leq nr_{j+1}$. Repeat the partitioning and limiting argument around (5.8)–(5.9) to conclude that $y^-(x_j, t) \leq r_{j+1}$, and then by definition $x^+(r_{j+1}, t) \geq x_j$. Now we have contradicted $x_j = x(r_j, t) + \varepsilon > x(r_{j+1}, t) + \varepsilon/2$.

It remains to show that assuming $X_n(nt) < nx(n^{-1}X_n(0), t) - n\varepsilon$ for infinitely many n also leads to a contradiction. Combine this with (5.16) to get

$$X_n(nt) < nx(n^{-1}X_n(0), t) - n\varepsilon \leq nx(r_{j+1}, t) - n\varepsilon \equiv nx'_j.$$

As in the proof of Proposition 5.2, switch to the \tilde{z}_n processes to get

$$\tilde{z}_n([nx'_j], nt) = \tilde{z}_n(k, 0) - \xi^k([nx'_j] - k, nt)$$

for some k such that $nr_j \leq X_n(0) \leq k \leq nc_1$, where c_1 is a fixed constant. From this conclude that $y^+(x'_j, t) \geq r_j$, which implies $x(r_j, t) = x^-(r_j, t) \leq x'_j = x(r_{j+1}, t) - \varepsilon$, a contradiction. \square

To prove Theorem 4(b), first a lemma

Lemma 5.2. *Let c be a finite constant.*

(a) *Suppose that $X_n(0) \geq nc$ for large enough n , almost surely. Then for any $x < c - t$, $X_n(nt) \geq nx$ for large enough n , almost surely.*

(b) *Suppose that $X_n(0) \leq nc$ for large enough n , almost surely. Then for any $x > c + t$, $X_n(nt) \leq nx$ for large enough n , almost surely.*

Proof. (a) Pick $\varepsilon > 0$ so that $x < c - t - 2\varepsilon$, and set $c_1 = c - \varepsilon$.

As an intermediate claim, we prove that almost surely, for large enough n ,

$$(5.17) \quad \xi^k([nx] - k, nt) = -K \cdot ([nx] - k) \quad \text{for all } k \geq [nx] + [n(t + \varepsilon)].$$

In other words, the claim is that $\xi^k([nx] - k, \cdot)$ has not jumped by time nt . The time of the first jump of $\xi^k([nx] - k, \cdot)$ is distributed as $\sum_{j=0}^{k-[nx]} Y_j$ where $\{Y_j\}$ are i.i.d. rate 1 exponential random variables. The estimate $P(\sum_{j=1}^N Y_j \leq Ns) \leq \exp[-N\kappa(s)]$ is valid for $0 < s < 1$. The rate function is $\kappa(s) = s - 1 - \log s$ for $s > 0$, which satisfies $\kappa'(s) < 0 < \kappa(s)$ for $0 < s < 1$. Thus

$$\begin{aligned} P\{(5.17) \text{ fails}\} &\leq \sum_{k \geq [nx] + [n(t+\varepsilon)]} P\left(\sum_{j=0}^{k-[nx]} Y_j \leq nt\right) \\ &\leq \sum_{i \geq [n(t+\varepsilon)]} \exp[-i\kappa(nt/i)] \leq C_1 e^{-C_2 n} \end{aligned}$$

for finite positive constants C_1, C_2 . This proves the claim, and (5.17) holds for large enough n a.s.

Since $\eta_n(X(0), 0) \leq K - 1$ and $[nc_1] < X(0)$,

$$z_n(k, 0) - z_n([nc_1], 0) = \sum_{i=[nc_1]+1}^k \eta_n(i, 0) \leq K(k - [nc_1]) - 1$$

for any $k \geq X_n(0)$. For the fixed x any $k \geq X_n(0)$ falls within the range in (5.17) for large enough n . So for large enough n ,

$$\begin{aligned} z_n(k, 0) - \xi^k([nx] - k, nt) &= z_n(k, 0) + K([nx] - k) \\ &\leq z_n([nc_1], 0) + K([nx] - [nc_1]) - 1 = z_n([nc_1], 0) - \xi^{[nc_1]}([nx] - [nc_1], nt) - 1. \end{aligned}$$

The last equality comes from (5.17) again because $[nc_1]$ also falls within the range of k 's in (5.17) for large enough n . The string of inequalities shows that no $k \geq X_n(0)$ can give the supremum in the microscopic variational formula (4.7) for $z_n([nx], nt)$, because each $k \geq X_n(0)$ is inferior to $[nc_1]$. But if $X_n(nt) < [nx]$, by (4.12) some $k \geq X_n(0)$ would be a maximizer. This contradiction shows that $X_n(nt) \geq [nx]$ for large enough n , a.s.

The proof of part (b) is similar, but instead of (5.17) use that $\xi^k([nx] - k, nt) = 0$ for all $k \leq [nx] - [n(t + \varepsilon)]$. \square

Proof of Theorem 4(b). The goal is to show that

$$(5.18) \quad \lim \left| \frac{X_n(nt)}{X_n(ns)} - x \left(\frac{X_n(ns)}{X_n(ns)}; s, t \right) \right| = 0$$

Let $\bar{u}_0(x) = u(x, s)$, $\bar{u}(x, \tau) = u(x, s + \tau)$ for $\tau > 0$, and let $\bar{x}(b; 0, \tau)$ denote characteristics for the solution \bar{u} . Then $x^\pm(b; s, t) = \bar{x}^\pm(b; 0, t - s)$. From the assumptions and Proposition 3.1 we know \bar{u} has unique characteristics: $\bar{x}(b; 0, \tau) = \bar{x}^\pm(b; 0, \tau)$.

Let $\bar{z}_n(i, t) = z_n(i, ns + t)$ be the n th process restarted at time ns . Let $\bar{X}_n(t)$ be the location of a second class particle in the process \bar{z}_n , with initial location $\bar{X}_n(0) = X_n(ns)$. In restarted terms, (5.18) is the same as

$$(5.19) \quad \lim_{n \rightarrow \infty} \left| \frac{\bar{X}_n(n(t-s))}{n} - \bar{x} \left(\frac{\bar{X}_n(0)}{n}; 0, t-s \right) \right| = 0 \quad \text{a.s.}$$

(5.19) follows from Theorem 4(a) if we check that the ingredients of the proof are valid for the processes \bar{z}_n and \bar{X}_n .

(i) We can obtain the variational formulas (4.7) and (4.9) for \bar{z}_n and \bar{X}_n in a form where the ξ -part depends on n through the time shift. For any fixed restarting time $\tau \geq 0$, let θ_τ denote the restarting operation on the probability space of the Poisson processes $\{D_i\}$: if the epochs of D_i in $(0, \infty)$ are $t_1 < t_2 < t_3 < \dots$ with $t_{j-1} \leq \tau < t_j$, then the epochs of $\theta_\tau D_i$ in $(0, \infty)$ are $t_j - \tau < t_{j+1} - \tau < t_{j+2} - \tau < \dots$.

The same proofs that originally gave Lemma 4.1 and Proposition 4.1 also yield, for $t > \tau$,

$$(5.20) \quad z(i, t) = \sup_k \{z(k, \tau) - \xi^k(i - k, t - \tau) \circ \theta_\tau\}$$

and

$$(5.21) \quad X(t) = \inf\{i : z(i, t) = z(k, \tau) - \xi^k(i - k, t - \tau) \circ \theta_\tau \text{ for some } k \geq X(\tau)\}.$$

Now for process z_n take $\tau = ns$, and (5.20)–(5.21) give for $t > 0$

$$\bar{z}_n(i, t) = \sup_k \{\bar{z}_n(k, 0) - \xi^k(i - k, t) \circ \theta_{ns}\}$$

and

$$\bar{X}_n(t) = \inf\{i : \bar{z}_n(i, t) = \bar{z}_n(k, 0) - \xi^k(i - k, t) \circ \theta_{ns} \text{ for some } k \geq X_n(ns)\}.$$

(ii) Limits (5.1) and (5.3) are true by definition of \bar{z}_n and \bar{u} . Limits (5.2) for $\xi^{[nr]}([nx], nt) \circ \theta_{ns}$ are true by the proofs in Seppäläinen (1999) because these limits are derived from summable deviation estimates and the Borel-Cantelli lemma. The distribution of $\xi^{[nr]}$ is not changed by the time shift, so the same deviation estimates are valid.

(iii) Finally assumption (2.25) for $\bar{X}_n(0) = X_n(ns)$ is a consequence of (2.25) for $X_n(0)$ and Lemma 5.2. \square

REFERENCES

- Dafermos, C. M. (1977). Generalized characteristics and the structure of solutions of hyperbolic conservation laws. *Indiana Univ. Math. J.* 26 1097–1119.
- L. C. Evans (1998). *Partial Differential Equations*. American Mathematical Society.
- Ferrari, P. A. (1992). Shock fluctuations in asymmetric simple exclusion. *Probab. Theory Related Fields* 91 81–101.
- P. A. Ferrari and L. R. G. Fontes (1994a). Shock fluctuations in the asymmetric simple exclusion process. *Probab. Theory Related Fields* 99 305–319.
- P. A. Ferrari and L. R. G. Fontes (1994b). Current fluctuations for the asymmetric simple exclusion process. *Ann. Probab.* 22 820–832.
- P. A. Ferrari, L. R. G. Fontes and Y. Kohayakawa (1994). Invariant measures for a two-species asymmetric process. *J. Statist. Phys.* 76 1153–1177.
- Ferrari, P. A. and Kipnis C. (1995). Second class particles in the rarefaction fan. *Ann. Inst. H. Poincaré Probab. Statist.* 31 143–154.
- H. Ishii (1984). Uniqueness of unbounded viscosity solutions of Hamilton-Jacobi equations. *Indiana Univ. Math. J.* 33 721–748.
- C. Kipnis and C. Landim (1999). *Scaling Limits of Interacting Particle Systems*. Grundlehren der mathematischen Wissenschaften, vol 320, Springer Verlag, Berlin.
- T. M. Liggett (1999). *Stochastic Interacting Systems: Contact, Voter and Exclusion Processes*. Springer-Verlag, New York.
- Rezakhanlou, F. (1995). Microscopic structure of shocks in one conservation laws. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 12 119–153.
- R. T. Rockafellar (1970). *Convex Analysis*. Princeton University Press.
- Rost, H. (1981). Non-equilibrium behaviour of a many particle process: Density profile and local equilibrium. *Z. Wahrsch. Verw. Gebiete* 58 41–53.
- T. Seppäläinen (1996). A microscopic model for the Burgers equation and longest increasing subsequences. *Electronic J. Probab.* 1, Paper 5, 1–51.
- T. Seppäläinen (1998a). Hydrodynamic scaling, convex duality, and asymptotic shapes of growth models. *Markov Process. Related Fields* 4 1–26.
- T. Seppäläinen (1998b). Coupling the totally asymmetric simple exclusion process with a moving interface. *Proceedings of I Escola Brasileira de Probabilidade (IMPA*

Rio de Janeiro, 1997). Markov Process. Related Fields 4 593–628.

T. Seppäläinen (1999). Existence of hydrodynamics for the totally asymmetric simple K -exclusion process. Ann. Probab. 27 361–415.

T. Seppäläinen (2000). A variational coupling for a totally asymmetric exclusion process with long jumps but no passing. To appear in the Proceedings of the Workshop on Hydrodynamic Limits, The Fields Institute, Toronto, 1998.